On the approximation properties of bivariate $(p,q)-$Bernstein operators (Revised)

Ali Karaisa*

Department of Mathematics–Computer Sciences, Faculty of Sciences, Necmettin Erbakan University, 42090 Meram, Konya, Turkey

arXiv:1601.05250v2 [math.CA] 2 Feb 2016

Abstract. In the present study, we have given a corrigendum to our paper on the approximation properties of bivariate $(p,q)-$Bernstein operators. Recently, we [2] have defined the bivariate $(p,q)-$Bernstein operators. Later, we have aware of Acar et. al [1] already have given some moments. In this case, we have revised [2, Lemma 2.3].

Keywords:bivariate $(p,q)-$**Bernstein operator, Voronovskaja type theorem,** $(p,q)-$**integer**
MSC:41A25, 41A36

1. Introduction

Approximation theory has been used in the theory of approximation of continuous functions by means of sequences of positive linear operators and still remains as a very active area of research. Since Korovkins famous theorem in 1950, the study of the linear methods of approximation given by sequences of positive and linear operators has became a firmly entrenched part of approximation theory.

During the last two decades, the applications of $q-$calculus have emerged as a new area in the field of approximation theory. The first $q-$analogue of the well-known Bernstein polynomials was introduced by Lupaş [12] by applying the idea of $q-$integers. Since approximation studied by $q-$Bernstein polynomials is better than classical one under convenient choice of q, many authors introduced q-generalization of various operators and investigated several approximation properties we refer the readers to [3–5].

Recently, Mursaleen et al. used (p,q)-calculus in approximation theory and defined $(p,q)-$analogue of Bernstein operators [13]. They estimated uniform convergence of the operators and rate of convergence, obtained Voronovskaya theorem as well. Also, $(p,q)-$analogue of Bernstein-Stancu operators and Bleimann-Butzer-Hahn operators were introduced in [14] and [15], respectively. For some recent works devoted to (p,q)-operators, we can refer the readers to [6–10].

In the present study, we define the bivariate Bernstein operators based on $(p,q)-$integer. We examine approximation properties of our new operator by the help of Korovkin-type theorem . Further, we present the local approximation properties and establish the rates of convergence by means of the modulus of continuity and the Lipschitz type maximal function. Also, we present a Voronovskaja type asymptotic formula for this operators.

Let us recall some definitions and notations regarding the concept of $(p,q)-$calculus.

The $(p,q)-$integer of the number n is defined by

$$[n]_{p,q} := \frac{p^n - q^n}{p-q}, \;\; n = 1,2,3\ldots, \;\; 0 < q < p \leqslant 1.$$

The $(p,q)-$factorial $[n]_{p,q}!$ and the $(p,q)-$binomial coefficients are defined as :

$$[n]_{p,q}! := \begin{cases} [n]_{p,q}\,[n-1]_{p,q}\cdots[1]_{p,q}\,, & n \in \mathbb{N} \\ 1, & n = 0 \end{cases}.$$

and

$$\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}!\,[n-k]_{p,q}!}, 0 \leqslant k \leqslant n.$$

Further, the $(p,q)-$binomial expansions are given as

$$(ax+by)^n_{p,q} = \sum_{k=0}^{n} p^{\binom{n-k}{2}} q^{\binom{k}{2}} a^{n-k} b^k x^{n-k} y^k.$$

and

$$(x-y)^n_{p,q} = (x-y)(px-qy)(p^2x-q^2y)\cdots(p^{n-1}x-q^{n-1}y).$$

Further information related to $(p,q)-$calculus can be found in [16, 17].

*Corresponding author: Tel:+90 332 323 8220; fax:+90 332 323 8245 E-mail:akaraisa@hotmail.com

2. Construction of the operators

Recently, Mursaleen et al applied $(p,q)-$calculus in approximation theory and introduced revised $(p,q)-$analogue of Bernstein operators as follows;

$$B_{n,p,q}(f;x)=\frac{1}{p^{n(n-1)/2}}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{k(k-1)/2}x^k\prod_{s=0}^{n-k-1}\left(p^s-q^s\frac{x}{b_n}\right)f\left(\frac{[k]_{p,q}}{[n]_{p,q}\,p^{k-n}}\right), \tag{2.1}$$

On the other hand, another active research area in approximation theory is to approximate the bivariate functions. For example, Barbosu, [19] defined and studied bivariate Bernstein operator. Büyükyazıcı[20] introduced $q-$Bernstein Chlodowsky operator.

Now, we define bivariate Bernstein operator based on $(p,q)-$integers. Let $I=[0,1]\times[0,1]$, $f:I\longrightarrow R$ and $0<q_1,q_2<p_1,p_2\leqslant 1$. We define the bivariate extension of the $(p,q)-$Bernstein operator operators as follows:

$$B_{n,m}^{(p_1,q_1),(p_1,q_1)}(f;x,y)=\sum_{k=0}^{n}\sum_{j=0}^{m}R_{n,k}(p_1,q_1;x)R_{m,j}(p_2,q_2;y)f\left(\frac{[k]_{p_1,q_1}}{[n]_{p_1,q_1}\,p_1^{k-n}},\frac{[j]_{p_2,q_2}}{[m]_{p_2,q_2}\,p_2^{j-m}}\right), \tag{2.2}$$

where

$$R_{n,k}(p_1,q_1;x)=p_1^{\frac{k(k-1)-n(n-1)}{2}}\left[\begin{array}{c} n \\ k \end{array}\right]_{p_1,q_1}x^k\prod_{s=0}^{n-k-1}(p_1^s-q_1^sx)$$

Lemma 2.1. *[11, Lemma 1]*

$$\begin{aligned} B_{n,p,q}\left(e^0;x\right) &= 1,\\ B_{n,p,q}\left(e_1;x\right) &= x,\\ B_{n,p,q}\left(e_2;x\right) &= \frac{p^{n-1}}{[n]_{p,q}}x+\frac{q[n-1]_{p,q}}{[n]_{p,q}}x^2. \end{aligned}$$

Also, $(p,q)-$Bernstein operator satisfy following equations:

Lemma 2.2.

$$\begin{aligned} B_{n,p,q}\left(e_3;x\right) &= \frac{p^{2n-2}}{[n]_{p,q}^2}x+\frac{p^{n-1}(2p+q)q[n-1]_{p,q}}{[n]_{p,q}^2}x^2+\frac{q^3[n-1]_{p,q}[n-2]_{p,q}}{[n]_{p,q}^2}x^3,\\ B_{n,p,q}\left(e_4;x\right) &= \frac{p^{3n-3}}{[n]_{p,q}^3}+\frac{q(3p^2+3qp+q^3)[n-1]_{p,q}p^{2n-4}}{[n]_{p,q}^3}x^2\\ &\quad+\frac{q^3(3p^2+2pq+q^2)[n-1]_{p,q}[n-2]_{p,q}p^{n-3}}{[n]_{p,q}^3}x^3+\frac{q^6[n-1]_{p,q}[n-2]_{p,q}[n-3]_{p,q}x^4}{[n]_{p,q}^3}, \end{aligned}$$

where $e_i(x)=x^i$, $i=0,1,2,3,4$.

Proof. Let us we compute e_3

$$\begin{aligned} B_{n,p,q}\left(e_3;x\right) &= \frac{1}{p^{n(n-7)/2}}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q}p^{k(k-7)/2}x^k\prod_{s=0}^{n-k-1}(p^s-q^sx)\frac{[k]_{p,q}^3}{[n]_{p,q}^3}\\ &= \frac{1}{p^{n(n-7)/2}\,[n]_{p,q}^2}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}p^{(k+1)(k-6)/2}[k+1]_{p,q}^2x^{k+1}\prod_{s=0}^{n-k-2}(p^s-q^sx). \end{aligned}$$

By $[k+1]_{p,q}^2 = p^{2k} + 2qp^k[k]_{p,q} + q^2[k]_{p,q}^2$, we have

$$\begin{aligned}
&= \frac{1}{p^{n(n-7)/2}[n]_{p,q}^2}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q} p^{(k^2-k-6)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&+2q\frac{1}{p^{n(n-7)/2}[n]_{p,q}^2}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}[k]_{p,q}p^{(k^2-3k-6)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&+\frac{q^2}{p^{n(n-7)/2}[n]_{p,q}^2}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}[k]_{p,q}^2p^{(k+1)(k-6)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&= \frac{x}{[n]_{p,q}^2}p^{2n-2} + \frac{2q[n-1]_{p,q}x^2}{[n]_{p,q}^2}p^{n-1} + \frac{p^{n-2}q^2[n-1]_{p,q}x^2}{[n]_{p,q}^2} + \frac{q^3[n-1]_{p,q}[n-2]_{p,q}x^3}{[n]_{p,q}^2}\\
&= \frac{x}{[n]_{p,q}^2}p^{2n-2} + \frac{(2p+q)q[n-1]_{p,q}x^2}{[n]_{p,q}^2}p^{n-2} + \frac{q^3[n-1]_{p,q}[n-2]_{p,q}x^3}{[n]_{p,q}^2}.
\end{aligned}$$

Finally,

$$\begin{aligned}
B_{n,p,q}(e_4;x) &= \frac{1}{p^{n(n-9)/2}}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q}x^k\prod_{s=0}^{n-k-1}(p^s - q^s x)\,p^{k(k-1)/2}p^{-4k}\frac{[k]_{p,q}^4}{[n]_{p,q}^4}\\
&= \frac{b_n^4}{p^{n(n-9)/2}[n]_{p,q}^3}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}p^{(k+1)(k-8)/2}[k+1]_{p,q}^3x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x).
\end{aligned}$$

Using the fact $[k+1]_{p,q}^3 = p^{3k} + 3p^{2k}q[k]_{p,q} + 3p^kq^2[k]_{p,q}^2 + q^3[k]_{p,q}^3$, we obtain

$$\begin{aligned}
B_{n,p,q}(e_4;x) &= \frac{p^{3n-3}}{p^{(n-1)(n-2)/2}[n]_{p,q}^3}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}p^{k(k-1)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&+\frac{3qb_n^4}{p^{n(n-9)/2}[n]_{p,q}^3}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}[k]_{p,q}p^{(k^2-3k-8)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&+\frac{3q^2}{p^{n(n-9)/2}[n]_{p,q}^3}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}[k]_{p,q}^2p^{(k^2-5k-8)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&+\frac{q^3}{p^{n(n-9)/2}[n]_{p,q}^3}\sum_{k=0}^{n-1}\left[\begin{array}{c} n-1 \\ k \end{array}\right]_{p,q}[k]_{p,q}^3p^{(k+1)(k-8)/2}x^{k+1}\prod_{s=0}^{n-k-2}(p^s - q^s x)\\
&= \frac{x}{[n]_{p,q}^3}p^{3n-3} + \frac{q(3p^2+3qp+q^2)[n-1]_{p,q}x^2}{[n]_{p,q}^3}p^{2n-4}\\
&+\frac{q^3(3p^2+2pq+q^2)[n-1]_{p,q}[n-2]_{p,q}x^3}{[n]_{p,q}^3}p^{n-3} + \frac{q^6[n-1]_{p,q}[n-2]_{p,q}[n-3]_{p,q}x^4}{[n]_{p,q}^3}.
\end{aligned}$$

Acar et. al [1] introduced Kantotovich modifications of $(p,q)-$Bernstein operators for bivariate functions using a new $(p,q)-$integral and given following moments for bivariate $(p,q)-$Bernstein operators.

Lemma 2.3. *[1, Lemma 1]*

$$\begin{aligned}
B_{n,m}^{(p_1,q_1),(p_2,q_2)}(1;x,y) &= 1,\\
B_{n,m}^{(p_1,q_1),(p_2,q_2)}(s;x,y) &= x,\\
B_{n,m}^{(p_1,q_1),(p_2,q_2)}(t;x,y) &= y,\\
B_{n,m}^{(p_1,q_1),(p_2,q_2)}(st;x,y) &= xy,\\
B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(s^2;x,y\right) &= \frac{p_1^{n-1}}{[n]_{p_1,q_1}}x + \frac{q_1[n-1]_{p_1,q_1}}{[n]_{p_1,q_1}}x^2,\\
B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(t^2;x,y\right) &= \frac{p_2^{m-1}}{[m]_{p_2,q_2}}y + \frac{q_2[m-1]_{p_2,q_2}}{[n]_{p_2,q_2}}y^2.
\end{aligned}$$

Using Lemma 2.3 and by linearity of $B_{n,m}^{(p_1,q_1),(p_2,q_2)}$, we have

Remark 2.4.

$$B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((t-x)^2;x,y\right) = \frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2), \tag{2.3}$$

$$B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s-y)^2;x,y\right) = \frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2). \tag{2.4}$$

The Korovkin-type theorem for functions of two variables was proved by Volkov [18].

Theorem 2.5. *Let $q_1 := (q_{1,n})$, $p_1 := (p_{1,n})$, $q_2 := (q_{m,2})$, $p_2 := (p_{m,2})$ such that $0 < q_{1,n}, q_{2,m} < p_{n,1}, p_{m,2} \leqslant 1$. If*

$$\lim_n p_{n,1} = 1,\ \lim_n q_{n,1} = 1,\ \lim_m p_{m,2} = 1,\ \lim_m q_{m,2} = 1,\ \lim_n p_{n,1}^n = a_1 \text{ and } \lim_m p_{m,1}^m = a_2, \tag{2.5}$$

the sequence $B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)$ convergence uniformly to $f(x,y)$, on $[0,1]\times[0,1] = [0,1]^2$ for each $f \in C\left([0,1]^2\right)$, where a_1, a_2 be reel numbers and $C([0,1]^2)$ be the space of all real valued continuous function on $[0,1]^2$ with the norm

$$\| f \|_{C([0,1]^2)} = \sup_{(x,y)\in[0,1]^2} |f(x,y)|.$$

Proof. Assume that the equities (2.5) are holds. Then, we have

$$\frac{p_{1,n}^{n-1}}{[n]_{p_{1,n},q_{1,n}}} \to 0,\ \frac{p_2^{m-1}}{[m]_{p_{2,m},q_{2,m}}} \to 0,\ \frac{q_{1,n}\,[n]_{p_{1,n},q_{1,n}}}{[n]_{p_{1,n},q_{1,n}}} \to 1 \text{ and } \frac{q_{2,m}\,[m-1]_{p_{2,m},q_{2,m}}}{[m]_{p_{2,m},q_{2,m}}} \to 1.$$

From Lemma 2.3, we obtain $\lim_{n,m\to\infty} B_{n,m}^{(p_1,q_1),(p_2,q_2)}(e_{ij};x,y) = e_{ij}(x,y)$ uniformly on $[0,1]^2$, where $e_{ij}(x,y) = x_i y_j, 0 \leqslant i+j \leqslant 2$ are the test functions. By using Korovkin theorem for functions of two variables was presented by Volkov [18], it follows that $\lim_{n,m\to\infty} B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y) = f(x,y)$, uniformly on $[0,1]^2$, for each $f \in C([0,1]^2)$. □

3. Rate of Convergence

In this section, we compute the rates of convergence of operators $B_{n,m}^{(p_1,q_1),(p_2,q_2)}$ to $f(x,y)$ by means of the modulus of continuity. Proceeding further, we provide a summary of the notations and definitions of the modulus of continuity and the Peetres $K-$functional for bivariate real valued functions.

For $f \in C([0,1]^2)$, the complete modulus of continuity for a bivariate case is defined as follows:

$$\omega(f,\delta) = \sup\left\{|f(t,s)-f(x,y)| : \sqrt{(t-x)^2+(s-y)^2} \leqslant \delta\right\}.$$

for every $(t,s),(x,y) \in [0,1]^2$. Further, partial moduli of continuity with respect to x and y are defined as

$$\begin{aligned}\omega^1(f,\delta) &= \sup\{|f(x_1,y)-f(x_2,y)| : y \in [0,1] \text{ and } |x_1-x_2| \leqslant \delta\}\\ \omega^2(f,\delta) &= \sup\{|f(x,y_1)-f(x,y_2)| : x \in [0,1] \text{ and } |y_1-y_2| \leqslant \delta\},\end{aligned}$$

It is obvious that they satisfy the properties of the usual modulus of continuity [21].

For $\delta > 0$, the Peetre-K functional [22] is given by

$$K(f,\delta) = \inf_{g\in C^2[0,1]^2}\left\{\|f-g\|_{C[0,1]^2} + \delta\|g\|_{C^2[0,1]^2}\right\},$$

where $C^2[0,1]^2$ is the space of functions of f such that f, $\frac{\partial^j f}{\partial x^j}$ and $\frac{\partial^j f}{\partial y^j}$ $(j=1,2)$ in $C[0,1]^2$. The norm $\|.\|$ on the space $C^2[0,1]^2$ is defined by

$$\|f\|_{C^2[0,1]^2} = \|f\|_{C[0,1]^2} + \sum_{j=1}^{2}\left(\left\|\frac{\partial^j f}{\partial y^j}\right\|_{C[0,1]^2} + \left\|\frac{\partial^j f}{\partial y^j}\right\|_{C[0,1]^2}\right).$$

Now, we give an estimate of the rate of convergence of operators $B_{n,m}^{(p_1,q_1),(p_2,q_2)}$.

Theorem 3.1. *Let $f \in C\left([0,1]^2\right)$. For all $x \in [0,1]^2$, we have*

$$\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)} - f(x,y)\right| \leqslant 2\omega\left(f;\delta_{n,m}\right),$$

where

$$\delta_{n,m}^2 = \frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2) + \frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2).$$

Proof. By definition the complete modulus of continuity of $f(x,y)$ and linearity and positivity our operator, we can write

$$\begin{aligned}|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)| &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|f(t,s)-f(x,y)|;x,y)\\ &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\omega\left(f;\sqrt{(t-x)^2+(s-y)^2}\right);x,y\right)\\ &\leqslant \omega(f,\delta_{n,m})\left[\frac{1}{\delta_{n,m}}B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\sqrt{(t-x)^2+(s-y)^2};x,y\right)\right].\end{aligned}$$

Using Cauchy-Scwartz inequality, from (2.3) and (2.4), one can write following

$$\begin{aligned}&|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)|\\ \leqslant\ &\omega(f,\delta_{n,m})\left[1+\frac{1}{\delta_{n,m}}\left\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((t-x)^2+(s-y)^2;x,y\right)\right\}^{1/2}\right]\\ =\ &\omega(f,\delta_{n,m})\left[1+\frac{1}{\delta_{n,m}}\left\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((t-x)^2;x,y\right)+B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s-y)^2;x,y\right)\right\}^{1/2}\right]\\ =\ &\omega(f,\delta_{n,m})\left[1+\frac{1}{\delta_{n,m}}\left(\frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2)+\frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2)\right)^{1/2}\right].\end{aligned}$$

Choosing $\delta_{n,m}=\left(\frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2)+\frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2)\right)^{1/2}$, for all $(x,y)\in[0,1]^2$, we get desired the result.

□

Theorem 3.2. *Let $f\in C\left([0,1]^2\right)$, then the following inequalities satisfy*

$$\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}-f(x,y)\right|\leqslant\omega^1(f;\delta_n)+\omega^2(f;\delta_m),$$

where

$$\delta_n^2 = \frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2), \tag{3.1}$$

$$\delta_m^2 = \frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2).. \tag{3.2}$$

Proof. By definition partial moduli of continuity of $f(x,y)$ and appliying Cauchy-Scwartz inequality, we have

$$\begin{aligned}|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)| &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|f(t,s)-f(x,y)|;x,y)\\ &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|f(t,s)-f(x,s)|;x,y)+B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|f(x,s)-f(x,y)|;x,y)\\ &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(|\omega^1(f;|t-x|)|;x,y\right)+B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(|\omega^2(f;|s-y|)|;x,y\right)\\ &\leqslant \omega^1(f,\delta_n)\left[1+\frac{1}{\delta_n}B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|t-x|;x,y)\right]\\ &\quad+\omega^2(f,\delta_m)\left[1+\frac{1}{\delta_m}B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|s-y|;x,y)\right]\\ &\leqslant \omega^1(f,\delta_n)\left[1+\frac{1}{\delta_n}\left(B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((t-x)^2;x,y\right)\right)^{1/2}\right]\\ &\quad+\omega^2(f,\delta_m)\left[1+\frac{1}{\delta_m}\left(B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s-y)^2;x,y\right)\right)^{1/2}\right].\end{aligned}$$

Consider (2.3), (2.4) and choosing

$$\begin{aligned}\delta_n^2 &= \frac{p_1^{n-1}}{[n]_{p_1,q_1}}(x-x^2),\\ \delta_m^2 &= \frac{p_2^{m-1}}{[m]_{p_2,q_2}}(y-y^2).\end{aligned}$$

we reach the result. □

For $\alpha_1,\alpha_1\in(0,1]$ and $(s,t),(x,y)\in[0,1]^2$, we define the Lipschitz class $LipM(\alpha_1,\alpha_1)$ for the bivariate case as follows:

$$|f(s,t)-f(x,y)|\leqslant M|s-x|^{\alpha_1}|t-y|^{\alpha_2}.$$

Theorem 3.3. *Let $f \in Lip_M(\alpha_1, \alpha_2)$. Then, for all $(x, y) \in [0, 1]^2$, we have*

$$|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| \leqslant M\delta_n^{\alpha_1/2}\delta_m^{\alpha_2/2},$$

where δ_n and δ_m defined in (3.1) and (3.2), respectively.

Proof. As $f \in Lip_M(\alpha_1, \alpha_2)$, it follows

$$\begin{aligned} |B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|f(t, s) - f(x, y)|, q_n; x, y) \\ &\leqslant MB_{n,m}^{(p_1,q_1),(p_2,q_2)}(|t - x|^{\alpha_1}|s - y|^{\alpha_2}; x, y) \\ &= MB_{n,m}^{(p_1,q_1),(p_2,q_2)}(|t - x|^{\alpha_1}|; x)B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|s - y|^{\alpha_2}; y). \end{aligned}$$

For $\widehat{p} = \frac{1}{\alpha_1}, \widehat{q} = \frac{\alpha_1}{2-\alpha_1}$ and $\widehat{p} = \frac{1}{\alpha_2}, \widehat{q} = \frac{\alpha_2}{2-\alpha_2}$ applying the Hölder's inequality, we get

$$\begin{aligned} |B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| &\leqslant M\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|t - x|^2; x)\}^{\alpha_1/2}\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(1; x)\}^{\alpha_1/2} \\ &\quad \times\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|s - y|^2; y)\}^{\alpha_{12}/2}\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(1; y)\}^{\alpha_2/2} \\ &= M\delta_n^{\alpha_1/2}\delta_m^{\alpha_2/2}. \end{aligned}$$

Hence, we get desired the result. □

Theorem 3.4. *Let $f \in C^1([0, 1]^2)$ and $0 < q_{1,n}, q_{2,m} < p_{n,1}, p_{m,2} \leqslant 1$. Then, we have*

$$|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| \leqslant \| f_x^{'} \|_{C([0,1]^2)} \delta_n + \| f_y^{'} \|_{C([0,1]^2)} \delta_m.$$

Proof. For $(t, s) \in [0, 1]^2$, we obtain

$$f(t) - f(s) = \int_x^t f_u^{'}(u, s)du + \int_y^s f_v^{'}(x, v)du$$

Applying the our operator on both sides above equation, we deduce

$$\begin{aligned} |B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| &\leqslant B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\left|\int_x^t f_u^{'}(u, s)du\right|; x, y\right) \\ &\quad + B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\left|\int_y^s f_v^{'}(x, v)du\right|; x, y\right). \end{aligned}$$

As

$$\left|\int_x^t f_u^{'}(u, s)du\right| \leqslant \| f_x^{'} \|_{C([0,1]^2)} |t - x| \text{ and } \left|\int_y^s f_v^{'}(x, v)du\right| \leqslant f_y^{'} \|_{C([0,1]^2)} |s - y|,$$

we have

$$\begin{aligned} |B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| &\leqslant f_x^{'} \|_{C([0,1]^2)} B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|t - x|; x, y) \\ &\quad + f_y^{'} \|_{C([0,1]^2)} B_{n,m}^{(p_1,q_1),(p_2,q_2)}(|s - y|; x, y). \end{aligned}$$

Using the Cauchy-Schwarz inequality, we can write following

$$\begin{aligned} |B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)| &\leqslant \| f_x^{'} \|_{C([0,1]^2)} \{B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((t - x)^2; x, y\right)\}^{1/2}\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(1; x, y)\}^{1/2} \\ &\quad + \| f_y^{'} \|_{C([0,1]^2)} \{B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s - y)^2; x, y\right)\}^{1/2}\{B_{n,m}^{(p_1,q_1),(p_2,q_2)}(1; x, y)\}^{1/2}. \end{aligned}$$

Form (2.3) and (2.4), we get desired the result. □

Theorem 3.5. *Let $f \in C\left([0, 1]^2\right)$, then we have*

$$\left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f; x, y) - f(x, y)\right\|_{C([0,1]^2)} \leqslant 2M\left(f; \delta_{n,m}(x, y)/2\right),$$

where

$$\delta_{n,m}(x, y) = \frac{1}{2}\max\left(\frac{p_1^{n-1}(x - x^2)}{[n]_{p_1,q_1}}, \frac{p_2^{m-1}(y - y^2}{[m]_{p_2,q_2}}\right).$$

Proof. Let $g \in C^2([0,1]^2)$. By the Taylor's formula, we get

$$\begin{aligned} g(s_1,s_2)-g(x,y) &= g(s_1,y)-g(x,y)+g(s_1,s_2)-g(s_1,y) \\ &= \frac{\partial g(x,y)}{\partial x}(s_1-x)+\int_x^{s_1}(s_1-u)\frac{\partial^2 g(u,y)}{\partial u^2}du \\ &\quad +\frac{\partial g(x,y)}{\partial x}(s_2-y)+\int_y^{s_2}(s_2-v)\frac{\partial^2 g(x,v)}{\partial v^2}dv \\ &= \frac{\partial g(x,y)}{\partial x}(s_1-x)+\int_0^{s_1-x}(s_1-x-u)\frac{\partial^2 g(u,y)}{\partial u^2}du \\ &\quad +\frac{\partial g(x,y)}{\partial x}(s_2-y)+\int_0^{s_2-y}(s_2-y-v)\frac{\partial^2 g(x,v)}{\partial v^2}dv \end{aligned}$$

Applying $B_{n,m}^{(p_1,q_1),(p_2,q_2)}$ to the both sides of the above equation, we obtain

$$\begin{aligned} \left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(g;x,y)-g(x,y)\right| &\leqslant \left|\frac{\partial g(x,y)}{\partial x}\right|\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}((s_1-x);x,y)\right| \\ &\quad +\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\int_0^{s_1-x}(s_1-x-u)\frac{\partial^2 g(u,y)}{\partial u^2}du;x,y\right)\right| \\ &\quad +\left|\frac{\partial g(x,y)}{\partial y}\right|\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}((s_2-y);x,y)\right| \\ &\quad +\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left(\int_0^{s_2-y}(s_2-y-v)\frac{\partial^2 g(v,x)}{\partial v^2}dv;x,y\right)\right| \end{aligned}$$

As $B_{n,m}^{(p_1,q_1),(p_2,q_2)}((s_1-x);x,y)=0$ and $B_{n,m}^{(p_1,q_1),(p_2,q_2)}((s_2-y);x,y)=0$, one can write following

$$\begin{aligned} \left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)\right\|_{C([0,1]^2)} &\leqslant \frac{1}{2}\left\|\frac{\partial g(x,y)}{\partial x}\right\|_{C([0,1]^2)}\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s_1-x)^2;x,y\right)\right| \\ &\quad +\frac{1}{2}\left\|\frac{\partial g(x,y)}{\partial y}\right\|_{C([0,1]^2)}\left|B_{n,m}^{(p_1,q_1),(p_2,q_2)}\left((s_2-y)^2;x,y\right)\right|. \end{aligned}$$

By (2.3), (2.4), we deduce,

$$\begin{aligned} \left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)\right\|_{C([0,1]^2)} &\leqslant \frac{1}{2}\max\left(\frac{p_1^{n-1}(x-x^2)}{[n]_{p_1,q_1}},\frac{p_2^{m-1}(y-y^2}{[m]_{p_2,q_2}}\right) \\ &\quad \times\left[\left\|\frac{\partial g(x,y)}{\partial x}\right\|_{C([0,1]^2)}+\left\|\frac{\partial g(x,y)}{\partial x}\right\|_{C([0,1]^2)}\right] \\ &\leqslant \|\,g\,\|_{C([0,1]^2)}\,\delta_{n,m}. \end{aligned} \tag{3.3}$$

By the linearity $B_{n,m}^{(p_1,q_1),(p_2,q_2)}$, we obtain

$$\begin{aligned} \left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)\right\|_{C([0,1]^2)} &\leqslant \left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}f-B_{n,m}^{(p_1,q_1),(p_2,q_2)}g\right\|_{C([0,1]^2)} \\ &\quad +\left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}g-g\right\|_{C([0,1]^2)}+\|f-g\|_{C([0,1]^2)}. \end{aligned} \tag{3.4}$$

By (3.5) and (3.6), one can see that

$$\left\|B_{n,m}^{(p_1,q_1),(p_2,q_2)}(f;x,y)-f(x,y)\right\|_{C([0,1]^2)} \leqslant 2M\left(f;\delta_{n,m}(x,y)/2\right).$$

This step completes the proof. □

First, we need the auxiliary result contained in the following lemma.

Lemma 3.6. *Let $0<q_n<p_n\leqslant 1$, be sequence such that $p_n,q_n\longrightarrow 1$ and $p_n^n\longrightarrow a$, $q_n^n\longrightarrow b$ as $n\longrightarrow\infty$. Then, we have the following limits:*

(i) $\lim_{n\to\infty}[n]_{q_n}B_n^{(p_n,q_n)}((t-x)^2;x)=ax-ax^2$

(ii) $\lim_{n\to\infty}[n]_{q_n}^2B_n^{(p_n,q_n)}((t-x)^4;x)=3ax^4-6ax^3+3ax^2$.

Proof. From (2.3), we get

$$[n]_{p_n,q_n}B_n^{(p_n,q_n)}((t-x)^2;x)=-p_n^{n-1}x^2+xp_n^{n-1}.$$

Let us take the limit of both sides of the above equality as $n \longrightarrow \infty$, then we have

$$\begin{aligned}\lim_{n\to\infty}[n]_{p_n,q_n}\left\{B_n^{(p_n,q_n)}((t-x)^2,x)\right\} &= \lim_{n\to\infty}\left\{-p_n^{n-1}x^2+xp_n^{n-1}\right\}\\ &= a(x-x^2).\end{aligned}$$

(ii) Using Lemma 2.1, Lemma 2.2 and by the linearity of the operators $B_n^{(p_n,q_n)}(f;x)$, we obtain

$$B_n^{(p_n,q_n)}(t-x)^4;x) = A_{1,n}x^4 + A_{2,n}x^3 + A_{3,n}x^2 + A_{4,n}x$$

where

$$\begin{aligned}A_{1,n} &= \frac{p_n^{n-3}[n]_{p_n,q_n}^2(-p_n^2+2p_nq_n-q_n^2)+p^{n-5}[n]_{p_n,q_n}(-p_n^3+3p_nq_n^2+q_n^3)-p_n^{3n-6}(p_n^2+p_n^3+2p_nq_n^2+q_n^3)}{[n]_{p_n,q_n}^3}\\ A_{2,n} &= \frac{p_n^{n-3}[n]_{p_n,q_n}^2(p_n^2-2p_nq_n+q_n^2)}{[n]_{p_n,q_n}^3}\\ &\quad+\frac{p^{2n-5}[n]_{p_n,q_n}(-q_n^3-4p_nq_n^2-3p_n^2q_n+2p_n^3)-p_n^{3n-6}(3p_n^3+3p_nq_n^2+5p_n^2q_n+q_n^3)}{[n]_{p_n,q_n}^3}\\ A_{3,n} &= \frac{p_n^{2n-4}[n]_{p_n,q_n}(-p_n^2+3p_nq_n+q_n^2)-p_n^{3n-5}(3p_n^2+q_n^2+3p_nq_n)}{[n]_{q_n}^3}\\ A_{4,n} &= \frac{p^{3n-3}}{[n]_{q_n}^3},\end{aligned}$$

It is clear that

$$\lim_{n\to\infty}[n]_{q_n}^2\{A_{4,n}x\}=0. \tag{3.5}$$

Taking the limit of both sides of $A_{1,n}$, we get

$$\begin{aligned}\lim_{n\to\infty}[n]_{q_n}^2\{A_{1,n}\} &= \lim_{n\to\infty}\left\{-p_n^{n-3}[n]_{p_n,q_n}(p_n-q_n)^2+p_n^{n-5}(-p_n^3+3p_nq_n^2+q_n^3)-\frac{p_n^{3n-6}(p_n^2+p_n^3+2p_nq_n^2+q_n^3)}{[n]_{q_n}}\right\}\\ &= \lim_{n\to\infty}\left\{-p_n^{n-3}(p_n^n-q_n^n)(p_n-q_n)+p_n^{n-5}(-p_n^3+3p_nq_n^2+q_n^3)-\frac{p_n^{3n-6}(p_n^2+p_n^3+2p_nq_n^2+q_n^3)}{[n]_{q_n}}\right\}\\ &= 3a. \qquad (3.6)\end{aligned}$$

Similarly, we can show that;

$$\lim_{n\to\infty}[n]_{q_n}^2\{A_{2,n}\}=-6a \text{ and } \lim_{n\to\infty}[n]_{q_n}^2\{A_{3,n}\}=3a. \tag{3.7}$$

By combining (3.5)-(3.7), we reach the desired the result. □

Now, we give a Voronovskaja type theorem for $B_{n,n}^{(p_n,q_n)}(f;x,y)$.

Theorem 3.7. *Let $f\in C^2([0,1]\times[0,1])$. Then, we have*

$$\lim_{n\to\infty}[n]_{p_n,q_n}B_{n,n}^{(p_n,q_n)}(f;x,y)-f(x,y)) = \frac{(ax-ax^2)f''_{x^2}(x,y)}{2}+\frac{(ay-ay^2)f''_{y^2}(x,y)}{2}.$$

Proof. Let $(x,y)\in[0,1]\times[0,1]$. Then, write Taylor's expansion of f as follows:

$$\begin{aligned}f(s,t) &= f(x,y)+f'_x(s-x)+f'_y(t-y)\\ &\quad+\frac{1}{2}\left\{f''_x(t-x)^2+2f'_{xy}(s-x)(t-y)+f''_y(t-y)^2\right\}+\varepsilon(s,t)\left((s-x)^2+(t-y)^2\right) \qquad (3.8)\end{aligned}$$

where $(s,t)\in[0,1]^2$ and $\varepsilon(s,t)\longrightarrow 0$ as $(s,t)\longrightarrow(x,y)$.

Applying the operator $B_{n,n}^{(p_n,q_n)}(f;.)$ on (3.8), we get

$$\begin{aligned}B_{n,n}^{(p_n,q_n)}(f;s,t)-f(x,y) &= f'_x(x,y)B_{n,n}^{(p_n,q_n)}((s-x);x,y)+f'_y(x,y)B_{n,n}^{(p_n,q_n)}((t-y);x,y)\\ &\quad+\frac{1}{2}\Bigg\{f''_{x^2}B_{n,n}^{(p_n,q_n)}((t-x_0)^2;x,y)+2f'_{xy}B_{n,n}^{(p_n,q_n)}((s-x)(t-y);x,y)\\ &\quad+f''_{y^2}B_{n,n}^{(p_n,q_n)}((t-y)^2;x,y)\Bigg\}+B_{n,n}^{(p_n,q_n)}\left(\varepsilon(s,t)\left((s-x)^2+(t-y)^2\right);x,y\right).\end{aligned}$$

Let us take the limit of both sides of the above equality as $n\longrightarrow\infty$, □

$$\lim_{n\to\infty}[n]_{p_n,q_n}B_{n,n}^{(p_n,q_n)}(f;s,t)-f(x,y)) = \lim_{n\to\infty}[n]_{p_n,q_n}\frac{1}{2}\Big\{f''_{x^2}B_{n,n}^{(p_n,q_n)}((t-x)^2;x,y) +2f'_{xy}B_{n,n}^{(p_n,q_n)}((s-x)(t-y);x,y)+f''_{y^2}B_{n,n}^{(p_n,q_n)}((t-y)^2;x,y)\Big\} + \lim_{n\to\infty}[n]_{p_n,q_n}B_{n,n}^{(p_n,q_n)}\left(\varepsilon(s,t)\left((s-x)^2+(t-y)^2\right);x,y\right).$$

For the last term on the right hand side, using Cauchy-Schwartz inequality, we get

$$B_{n,n}^{(p_n,q_n)}\left(\varepsilon(s,t)\left((s-x)^2+(t-y)^2\right);x,y\right) \leqslant \sqrt{\lim_{n\to\infty}B_{n,n}^{(p_n,q_n)}\left(\varepsilon^2(s,t);x,y\right)} \times \sqrt{2\lim_{n\to\infty}[n]_{p_n,q_n}^2B_{n,n}^{(p_n,q_n)}\left(\varepsilon(s,t)\left((s-x)^4+(t-y)^4\right);x,y\right)}.$$

As $\lim_{n\to\infty}B_{n,n}^{(p_n,q_n)}\left(\varepsilon^2(s,t);x,y\right)=\varepsilon^2(x,y)=0$ and using Lemma 3.6(ii)
$\lim_{n\to\infty}[n]_{p_n,q_n}^2B_{n,n}^{(p_n,q_n)}\left((s-x)^4+(t-y)^4);x,y\right)$ is finite, then we obtain

$$\lim_{n\to\infty}[n]_{p_n,q_n}^2B_{n,n}^{(p_n,q_n)}\left(\varepsilon(s,t)\left((s-x)^4+(t-y)^4\right);x,y\right)=0.$$

Hence, one can see that

$$\lim_{n\to\infty}[n]_{p_n,q_n}B_{n,n}^{(p_n,q_n)}(f;x,y)-f(x,y)) = \frac{(ax-ax^2)f''_{x^2}(x,y)}{2}+\frac{(ay-ay^2)f''_{y^2}(x,y)}{2}.$$

This step completes the proof.

□

References

[1] T. Acar, A. Aral, and S.A. Mohiuddine, approximation by bivariate $(p,q)-$Bernstein–Kantorovich operator arXiv:submit/1454072 [math.CA] 13 Jan 2016.

[2] A. Karaisa,On the approximation properties of bivariate $(p,q)-$Bernstein operators arXiv:1601.05250 [math.CA]

[3] A. Karaisa, D T. Tollu and Y.Asar, Stancu type generalization of q-Favard-Szàsz operators, Appl. Math. Comput. 264 (2015) 249–257.

[4] A. M. Acu, C. V. Muraru, Approximation Properties of bivariate extension of $q-$Bernstein-Schurer-Kantorovich operators, Result Math. 67 (3), (2015), 265–279.

[5] A. Aral, V.Gupta and Ravi P. Agarwal, Applications of $q-$calculus in operator theory. Berlin: Springer, 2013.

[6] V. Gupta, $(p,q)-$Szàsz-Mirakyan-Baskakov operators, Complex Analysis and Operator Theory, 1–9.

[7] T. Acar, $(p,q)-$generalization of Szàsz–Mirakyan operators, Math. Method. Appl.Sci., DOI: 10.1002/mma.3721.

[8] T. Acar, S. A. Mohiudine, M. Mursaleen, Approximation by $(p,q)-$Baskakov-Durrmeyer–Stancu operators, Communicated.

[9] M. Mursaleen, Md. Nasiuzzaman and A. Nurgali, Some approximation results on BernsteinSchurer operators defined by (p,q)-integers, Jou. Ineq. Appl., 2015 (2015) 249.

[10] A. Aral, V. Gupta, Bernstein Durrmeyer operators based on two parameters, Communicated.

[11] Mursaleen, M and Ansari, Khursheed J and Khan, Asif Some Approximation results by $(p,q)-$analogue of Bernstein–Stancu operators (Revised).

[12] A. Lupaş, $q-$analogue of the Bernstein operator, Seminar on Numerical and Statistical Calculus, University of Cluj-Napoca, 9 (1987), 85-92.

[13] M. Mursaleen, K. J. Ansari and A. Khan, On $(p,q)-$analogue of Bernstein operators, Appl. Math. Comput., 10.1016/j.amc.2015.04.090.

[14] M. Mursaleen, K. J. Ansari and A. Khan, Some approximation results by $(p,q)-$analogue of Bernstein–Stancu operators, Appl. Math. Comput., 10.1016/j.amc.2015.03.135.

[15] M. Mursaleen, Md. Nasiruzzaman, Asif Khan, Khursheed J. Ansari, Some approximation results on Bleimann-Butzer–Hahn operators defined by $(p,q)-$integers, arXiv:1505.00392, (2015).

[16] P. N. Sadjang, On the fundamental theorem of $(p,q)-$calculus and some $(p,q)-$Taylor formulas, arXiv:1309.3934 [math.QA].

[17] V. Sahai, S. Yadav, Representations of two parameter quantum algebras and $p,q-$special functions, J. Math. Anal. Appl., 335 (2007), 268279.

[18] V. J.Volkov, On the convergence of linear positive operators in the space of continuous functions of two variables (Russian), Dokl. akad. Nauk. SSSR (N.S), 115, 1957, 17-19.
[19] D. Barbosu , Some Generalized Bivariate Bernstein Operators, Mathematical Notes, Miskolc, 1 (1), (2000), 3–10.
[20] I. Büyükyazıcı, On the approximation properties of two-dimensional $q-$Bernstein-Chlodowsky polynomials, Math. Commun., 14(2) (2009),255–269.
[21] G. A. Anastassiou, S. G. Gal, Approximation theory: moduli of continuity and global smoothness preservation, Birkhäuser, Boston, 2000.
[22] P. L. Butzer, H. Berens, Semi-groups of operators and approximation, Springer, New York, 1967